

TECHNICAL DATA SHEET

CyFlow™ CD33 PE Anti-Hu; Clone WM53

REF BW164790



Sysmex Partec GmbH

Arndtstraße 11 a-b

02826 Görlitz

Tel +49 3581 8746 0

E-mail: info@sysmex-partec.com

www.sysmex-partec.com

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD33
Alternative Names	p67, SIGLECT-3
Clone	WM53
Clonality	monoclonal
Format	PE
Host / Isotype	Mouse / IgG1
Species Reactivity	Human, Non-Human Primates
Negative Species Reactivity	—
Quantity	100 tests
Immunogen	Human Acute Myelogenous Leukemia (AML) cells

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services



Specificity

The mouse monoclonal antibody WM53 recognize CD33 antigen, a 67 kDa type I transmembrane glycoprotein (immunoglobulin superfamily) expressed on myeloid progenitors, monocytes, granulocytes, dendritic cells and mast cells; it is absent on platelets, lymphocytes, erythrocytes and hematopoietic stem cells.

HLDA IV—WS Code M-505

Application

The reagent is designed for Flow Cytometry analysis of human blood cells. Recommended usage is 10- μ l reagent-/ 100- μ l of whole blood or 10^6 cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH \approx 7.4, containing 0.09% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD33 is a transmembrane protein of the sialic acid-binding immunoglobulin-like lectin (Siglec) family. It belongs to the immunoreceptor tyrosine-based inhibitory motif (ITIM)-containing molecules able of recruiting protein tyrosine phosphatases SHP-1 and SHP-2 to signal assemblies; these ITIMs are also used for ubiquitin-mediated removal of the receptor from the cell surface. CD33 is expressed on cells of myelomonocytic lineage, binds sialic acid residues in N- and O-glycans on cell surfaces, and is a therapeutic target for acute myeloid leukemia.

Warnings

Non-Hazardous Statement: This is not considered hazardous by the criteria in 29 CFR 1910.1200 or the General Classification guideline for preparations of the EU.

Safety Data Sheet Statement: Important information regarding the safe handling, transport, and disposal of this product is contained in the Safety Data Sheet.










The Safety Data Sheet for this product is available at www.sysmex-partec.com/services

References

- Favaloro EJ, Bradstock KF, Kabral A, Grimsley P, Berndt MC: Characterization of monoclonal antibodies to the human myeloid-differentiation antigen, 'gp67' (CD-33). *Dis Markers*. 1987 Dec; 5(4):215-25. < PMID: 3502990 >
- Favaloro EJ, Bradstock KF, Kabral A, Grimsley P, Zowtyj H, Zola H: Further characterization of human myeloid antigens (gp160, 95, gp150, gp67) investigation of epitopic heterogeneity and non-haemopoietic distribution using panels of monoclonal antibodies belonging to CD-11b, CD-13 and CD-33. *Br J Haematol*. 1988 Jun; 69(2):163-71. < PMID: 2455535 >
- Bradstock KF, Kirk J, Grimsley PG, Kabral A, Hughes WG: Unusual immunophenotypes in acute leukaemias: incidence and clinical correlations. *Br J Haematol*. 1989 Aug; 72(4):512-8. < PMID: 2673329 >
- Shin YK, Choi EY, Kim SH, Chung J, Chung DH, Park WS, Jung KC, Kim HS, Park S, Kim HJ, Park MH, Min CK, Kim CC, Park SH: Expression of leukemia-associated antigen, JL1, in bone marrow and thymus. *Am J Pathol*. 2001 Apr; 158(4):1473-80. < PMID: 11290565 >
- Vitale C, Romagnani C, Puccetti A, Olive D, Costello R, Chiossone L, Pitto A, Bacigalupo A, Moretta L, Mingari MC: Surface expression and function of p75/AIRM-1 or CD33 in acute myeloid leukemias: engagement of CD33 induces apoptosis of leukemic cells. *Proc Natl Acad Sci USA*. 2001 May 8; 98(10):5764-9. < PMID: 11320212 >
- Hernández-López C, Varas A, Sacedón R, Jiménez E, Muñoz JJ, Zapata AG, Vicente A: Stromal cell-derived factor 1/CXCR4 signaling is critical for early human T-cell development. *Blood*. 2002 Jan 15; 99(2):546-54. < PMID: 11781237 >
- Schenk M, Bouchon A, Birrer S, Colonna M, Mueller C: Macrophages expressing triggering receptor expressed on myeloid cells-1 are underrepresented in the human intestine. *J Immunol*. 2005 Jan 1; 174(1):517-24. < PMID: 15611278 >
- Garnache-Ottou F, Chaperot L, Biichle S, Ferrand C, Remy-Martin JP, Deconinck E, de Tailly PD, Bulabois B, Poulet J, Kuhlein E, Jacob MC, Salaun V, Arock M, Drenou B, Schillinger F, Seilles E, Tiberghien P, Bensa JC, Plumas J, Saas P: Expression of the myeloid-associated marker CD33 is not an exclusive factor for leukemic plasmacytoid dendritic cells. *Blood*. 2005 Feb 1; 105(3):1256-64. < PMID: 15388576 >
- Leone AM, Rutella S, Bonanno G, Abbate A, Rebuffi AG, Giovannini S, Lombardi M, Galiuto L, Liuzzo G, Andreotti F, Lanza GA, Contemi AM, Leone G, Crea F: Mobilization of bone marrow-derived stem cells after myocardial infarction and left ventricular function. *Eur Heart J*. 2005 Jun; 26(12):1196-204. < PMID: 15734770 >

- Hernández-Caselles T, Martínez-Esparza M, Pérez-Oliva AB, Quintanilla-Cecconi AM, García-Alonso A, Alvarez-López DM, García-Peñarrubia P: A study of CD33 (SIGLEC-3) antigen expression and function on activated human T and NK cells: two isoforms of CD33 are generated by alternative splicing. J Leukoc Biol. 2006 Jan; 79(1):46-58. < PMID: 16380601 >

Symbols

	Reagent catalogue number		Contains sufficient for <n> tests
	For Research Use Only. Not for use in diagnostic and therapeutic procedures.		Temperature limitation
	Batch code		Keep away from sunlight
	Manufacturer		Consult accompanying documents
	Use by		